

RAW Assessment – Clinician Guide

This guide explains the purpose and use of the RAW (Ready, Able, Willing) model in clinical settings. It is designed to support clinicians in evaluating a client's readiness for recovery and tailoring interventions accordingly.

Overview of RAW

The RAW framework helps clinicians answer:

- **READY** – Does the client recognize a need for change?
- **ABLE** – Do they have the tools and support to act?
- **WILLING** – Are they motivated to begin recovery work?

Each dimension is rated on a scale of 0–4, leading to a total readiness score of 0–12.

1. READY – Recognition of Need

Indicators of high readiness:

- Expresses dissatisfaction with current life
- Identifies personal goals or values being compromised
- Acknowledges risk or harm from continued use

Clinical Notes:

- Low scores may indicate denial or limited insight
- Motivational interviewing is useful to deepen awareness

2. ABLE – Capacity and Support

Indicators of strong ability:

- Stable housing, physical health, transportation

- Access to services, skills for emotion regulation
- Cognitive capacity to participate in recovery work

Clinical Notes:

- Scores of 0–1 suggest need for stabilization before deeper work
- Coordinate referrals for support services as needed

3. WILLING – Motivation and Engagement

Indicators of strong willingness:

- Self-initiated efforts or follow-through
- Verbal commitment to recovery
- Willingness to explore pain, values, identity

Clinical Notes:

- Resistance is not failure: explore ambivalence gently
- Celebrate micro-commitments, even symbolic ones

Using RAW in Practice

- Conduct RAW assessments during intake, reassessment, or case review
- Use results to shape pace, intensity, and type of intervention
- Repeat the assessment to track progress over time
- Combine with STAR Matrix for full complexity + readiness mapping